

PAPER • OPEN ACCESS

How many mutually unbiased bases are needed to detect bound entangled states?

To cite this article: Joonwoo Bae *et al* 2022 *J. Phys. A: Math. Theor.* **55** 505303

View the [article online](#) for updates and enhancements.

You may also like

- [Detecting entanglement can be more effective with inequivalent mutually unbiased bases](#)
B C Hiesmayr, D McNulty, S Baek et al.
- [Incompatibility robustness of quantum measurements: a unified framework](#)
Sébastien Designolle, Máté Farkas and Jdrzej Kaniewski
- [Orbits of mutually unbiased bases](#)
Kate Blanchfield

How many mutually unbiased bases are needed to detect bound entangled states?

Joonwoo Bae¹ , Anindita Bera² , Dariusz Chruściński² ,
Beatrix C Hiesmayr³  and Daniel McNulty^{4,5,*} 

¹ School of Electrical Engineering, Korea Advanced Institute of Science and Technology (KAIST), 291 Daehak-ro Yuseong-gu, Daejeon 34141, Republic of Korea

² Institute of Physics, Faculty of Physics, Astronomy and Informatics Nicolaus Copernicus University, Grudziądzka 5/7, 87–100 Toruń, Poland

³ Faculty of Physics, University of Vienna, Währinger Straße 17, 1090 Vienna, Austria

⁴ Center for Theoretical Physics, Polish Academy of Sciences, Al. Lotników 32/46, 02-668 Warsaw, Poland

⁵ Department of Mathematics, Aberystwyth University, Aberystwyth, Wales, United Kingdom

E-mail: dmcnulty@cft.edu.pl

Received 1 August 2022; revised 29 November 2022

Accepted for publication 8 December 2022

Published 28 December 2022



CrossMark

Abstract

From a practical perspective it is advantageous to develop methods that verify entanglement in quantum states with as few measurements as possible. In this paper we investigate the minimal number of mutually unbiased bases (MUBs) needed to detect bound entanglement in bipartite $(d \times d)$ -dimensional states, i.e. entangled states that are positive under partial transposition. In particular, we show that a class of entanglement witnesses (EWs) composed of MUBs can detect bound entanglement if the number of measurements is greater than $d/2 + 1$. This is a substantial improvement over other detection methods, requiring significantly fewer resources than either full quantum state tomography or measuring a complete set of $d + 1$ MUBs. Our approach is based on a partial characterisation of the (non-)decomposability of EWs. We show that non-decomposability is a universal property of MUBs, which holds regardless of the choice of complementary observables, and we find that both the number

* Author to whom any correspondence should be addressed.



Original Content from this work may be used under the terms of the [Creative Commons Attribution 4.0 licence](https://creativecommons.org/licenses/by/4.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

of measurements and the structure of the witness play an important role in the detection of bound entanglement.

Keywords: entanglement detection, bound entanglement, mutually unbiased bases, non-decomposable witnesses

1. Introduction

The extensive study of correlations in quantum systems has revealed a complex hierarchy of entanglement, each class with its own distinct non-classical properties and applications. Examples include entangled states which are unsteerable (admitting a local hidden state model [1]), steerable states (which admit only a local hidden variable model [2]), and entangled states which exhibit Bell non-locality [3]. There is also a class of states, known as bound entangled states, from which no pure entangled state can be distilled [4, 5]. Any entangled state that remains positive under partial transposition (PPT) is bound entangled, but it is not yet known if all bound entangled states are PPT.

Entanglement is a fundamental ingredient in almost all quantum information tasks. Usually, more entanglement in a system implies a higher degree of performance for certain tasks [6–8] and, consequently, highly entangled states are a valuable resource for quantum technologies. While bound entangled states are considered weakly entangled, they can be used for quantum steering (i.e. ruling out local hidden state models [9]), the distillation of secure quantum keys [10, 11], and for quantum teleportation [12]. They also provide valuable insights into the study of classical information theory, providing important connections to bound information [13].

Certifying non-classical correlations is both of fundamental and practical interest [14–16], and has been observed at low and high energies (e.g. in the entangled K-meson system [17], photons beyond the optical wave length [18], or oscillating neutrino systems [19]).

Given the full information about a quantum system, it is, in general, computationally hard to reveal entanglement [20]. More precisely, the difficulty lies in the characterisation of those entangled states that remain PPT. So far, only a few families of bound entangled states have been constructed [21–30], each by very different tools, and a general characterisation of bound entangled states is missing.

There are several criteria which enable the detection of entangled states (see [14–16]). A popular detection scheme is based on the concept of an entanglement witness (EW), a Hermitian operator \mathbf{W} that satisfies

$$\begin{aligned} \text{tr}[\mathbf{W}\rho_{\text{sep}}] &\geq 0 \text{ for all separable states } \rho_{\text{sep}}, \\ \text{and } \text{tr}[\mathbf{W}\rho] &< 0 \text{ for some entangled states } \rho. \end{aligned} \quad (1)$$

Hierarchical structures on the level of certifications of entanglement have been recently introduced according to the assumptions made in concrete experimental setups [31, 32]. In the *standard* scenario all experimental devices are trusted, whereas another extreme scenario is a device-independent one, which demands that the certification does not rely on the details of the source and the measurement devices [33]. An intermediate scenario is measurement-device-independent certification that is supposed to close all loopholes that may appear in the detection devices. In this case, the certification can be straightforwardly constructed from EWs [34, 35], which we pursue in this contribution.

Note that EWs correspond to a particular set of observables that satisfy equation (1). They can be decomposed into local measurements, where it is of practical importance to minimise the number of outcomes, or precisely, positive-operator-valued-measure (POVM) elements. An interesting class of EWs arises from sets of mutually unbiased bases (MUBs), that are

experimentally the most feasible and also gradually approach quantum state tomography as they converge to being complete.

A pair of orthonormal bases \mathcal{B}_1 and \mathcal{B}_2 of the space \mathbb{C}^d are mutually unbiased if the basis vectors satisfy $|\langle \psi_i | \varphi_j \rangle|^2 = \frac{1}{d}$ for all $|\psi_i\rangle \in \mathcal{B}_1$ and $|\varphi_j\rangle \in \mathcal{B}_2$. This condition mathematically formalises the concept of complementarity, i.e. if the system resides in an eigenstate of one observable, the outcome of measuring a second complementary observable is maximally uncertain. Generalising to larger collections of measurements, we say a set of bases is mutually unbiased if all bases are pairwise unbiased. As is well known (see [36] for a review), complete sets of $d + 1$ MUBs have been constructed in prime and prime-power dimensions [37, 38], but it is conjectured that only smaller sets exist in all other cases [39].

The first direct connection between MUBs and EWs was made in [40], with the construction of witnesses composed of two or more MUBs. Generalisations of the witnesses to a wider class of measurements, including 2-designs and mutually unbiased measurements, for which MUBs are an example, have since appeared [41–48]. In low dimensions, equipped with a complete set of MUBs, examples of these witnesses have been shown to experimentally detect bound entangled states [49]. However, little is known about their capability to detect PPT entangled states in arbitrary dimensions and the number of measurements required.

The ability of an EW to detect bound entangled states is directly connected to the notion of non-decomposability [5, 50–53]. An EW \mathbf{W} is decomposable if $\mathbf{W} = A + B^\Gamma$, with $A, B \geq 0$ (Γ denotes the partial transposition). EWs that cannot be decomposed in this way (i.e. non-decomposable) are able to detect bound entangled states. In fact, an EW is non-decomposable if and only if it detects PPT entangled states [51]. Unfortunately, there is no general method to construct such objects, and it is usually difficult to decide whether a witness is decomposable or not. One focus of this paper will be to partially ameliorate this difficulty by constructing a general class of non-decomposable witnesses from small sets of MUBs.

Another popular entanglement detection approach implements tomographically complete measurements to recover all information about the given quantum state. With full knowledge of the state, any known detection tool can then be applied to certify entanglement. In this scenario, complete sets of MUBs also appear, as they provide an optimal measurement scheme for quantum state tomography [37, 38]. For example, bound entanglement detection via quantum state tomography has been experimentally implemented several times before in the multipartite setting [54–58], and less frequently in the bipartite scenario [49, 59].

The two certification methods described above and their connection with MUBs is particularly convenient due to the relatively simple implementation of these observables experimentally. For example, experimental implementations of MUBs have been demonstrated in a wide range of quantum information protocols, e.g. [49, 60–68], as well as in fairly large dimensional systems [69, 70]. The EW approach has the added bonus of requiring at most $d + 1$ MUBs, measured on each subsystem, while full quantum state reconstruction requires $d^2 + 1$ MUBs in $\mathbb{C}^d \otimes \mathbb{C}^d$ (or any other informationally complete set of measurements on the full state space), and often suffers from errors, such as the stability of the source over all possible measurements [71, 72]. In general dimensions (not restricted to prime powers) one can use weighted two-designs consisting of orthonormal bases to perform quantum state tomography but this requires more bases than MUBs [73]. Known witnesses for bound entanglement, e.g. the computable cross-norm or realignment criterion [74], similarly require of the order d^2 measurements. Methods with as few measurements as possible are therefore highly desirable.

In this work, our primary motivation is to ask whether a complete set of MUBs in \mathbb{C}^d is needed to detect bound entangled states. First, we consider which of the previously constructed EWs, composed of a complete set of MUBs, are useful to detect bound entanglement. We

show (section 3) that the original witness described in [40] is decomposable for any d , and fails to detect PPT entangled states. We then demonstrate (section 4) that modifying the EW (via re-orderings of a single basis in one subsystem [43, 49]), results in a non-decomposable witness that detects PPT entangled states in any dimension d , provided a complete set of MUBs exists. We derive our main result in section 5, which shows that bound entanglement is detected by *any* set of m MUBs (in every dimension d) when $m > d/2 + 1$. This requires significantly fewer measurements than previous approaches. In section 6, we show that a pair of Fourier connected MUBs is incapable of certifying bound entanglement. We then conclude with examples in $d = 3$ (section 7) and $d = 2^r$ (section 8). In particular, we show that three MUBs are both necessary and sufficient to detect bound entanglement when $d = 3$. We conjecture that bound entanglement can be verified with only $m = d/2 + 1$ MUBs when $d = 2^r$, $r \geq 2$. This is corroborated with examples, in $d = 4$, of non-decomposable witnesses using only three measurements, and as a by-product we present new families of bound entangled states that they detect.

Overall, our work illustrates that MUBs are a powerful tool in the study of quantum correlations, useful not only to distinguish between separable and entangled states, but also to penetrate deeper into the full hierarchy of non-classical correlations. This contribution, together with recent work connecting MUBs with Bell nonlocality [75] and quantum steering [76], is further evidence of the fundamental and practical importance of this special class of measurements in revealing non-classical correlations.

2. EWs with MUBs

Let us start with the framework of EWs presented in [77], where it is clear that EWs are built from a collection of measurements. Suppose that a non-negative operator C is decomposed as

$$C = \sum_{a,b} A_a \otimes B_b,$$

where $\{A_a\}$ and $\{B_b\}$ POVM elements and refer to the two measurable subspaces of the quantum system of interest. There are non-trivial upper and lower bounds satisfied by all separable states ρ_{sep} , i.e.

$$L(C) \leq \text{tr}[\rho_{\text{sep}} C] \leq U(C).$$

Each of the bounds corresponds to an EW. Since upper bounds are useful to detect PPT entangled states [43, 49], we consider EWs that correspond to upper bounds. That is, a witness \mathbf{W} is constructed such that

$$\text{tr}[\mathbf{W}\rho] \not\leq 0 \iff \text{tr}[C\rho] \not\leq U(C),$$

for entangled states ρ detected by an upper bound.

In what follows, we derive EWs from a collection of MUBs. Let us consider m MUBs in a d -dimensional Hilbert space,

$$\mathcal{M}_m = \{\mathcal{B}_0, \mathcal{B}_1, \dots, \mathcal{B}_{m-1}\},$$

where $\mathcal{B}_\alpha = \{|i_\alpha\rangle\}_{i=0}^{d-1}$, and \mathcal{B}_0 is chosen as the canonical basis $\{|0\rangle, \dots, |d-1\rangle\}$. From these MUBs we construct the following operators

$$\mathbf{B}(\mathcal{M}_m, s) = \sum_{\ell=0}^{d-1} |\ell\rangle\langle\ell| \otimes |\ell+s\rangle\langle\ell+s| + \sum_{\alpha=1}^{m-1} \sum_{i=0}^{d-1} |i_\alpha\rangle\langle i_\alpha| \otimes |i_\alpha\rangle\langle i_\alpha|,$$

and its partial transpose

$$\mathbf{B}^\Gamma(\mathcal{M}_m, s) = \sum_{\ell=0}^{d-1} |\ell\rangle\langle\ell| \otimes |\ell+s\rangle\langle\ell+s| + \sum_{\alpha=1}^{m-1} \sum_{i=0}^{d-1} |i_\alpha\rangle\langle i_\alpha| \otimes |i_\alpha^*\rangle\langle i_\alpha^*|,$$

where the complex conjugation (and transposition) is performed with respect to the canonical basis \mathcal{B}_0 , and the sum $\ell + s$ is defined modulo d . It has been proven [40] that for any separable state, the two operators are bounded from above by the same quantity, depending on the number of MUBs, m , and the dimension d . In particular,

$$\text{tr}[\mathbf{B}(\mathcal{M}_m, s) \rho_{\text{sep}}] \leq \frac{d+m-1}{d}, \tag{2}$$

and

$$\text{tr}[\mathbf{B}^\Gamma(\mathcal{M}_m, s) \rho_{\text{sep}}] \leq \frac{d+m-1}{d}. \tag{3}$$

Hence, the operator

$$\mathbf{W}(\mathcal{M}_m, s) = \frac{d+m-1}{d} \mathbb{1}_d \otimes \mathbb{1}_d - \mathbf{B}(\mathcal{M}_m, s), \tag{4}$$

with $\mathbb{1}_d$ the $d \times d$ identity matrix, satisfies $\text{tr}[\mathbf{W}(\mathcal{M}_m, s) \rho_{\text{sep}}] \geq 0$. Similarly, $\text{tr}[\mathbf{W}^\Gamma(\mathcal{M}_m, s) \rho_{\text{sep}}] \geq 0$. It is, therefore, clear that whenever $\mathbf{W}(\mathcal{M}_m, s)$ possesses a strictly negative eigenvalue it defines an EW.

3. Witnesses with $s = 0$ do not detect bound entanglement

Consider first the simplest scenario $s = 0$, such that $\mathbf{W}^\Gamma(\mathcal{M}_m, 0)$ corresponds to the witness first considered in [40]. Suppose that the d -dimensional Hilbert space $\mathcal{H} = \mathbb{C}^d$ admits a complete set of $d + 1$ MUBs. The witness of equation (4) is therefore

$$\mathbf{W}(\mathcal{M}_{d+1}, 0) = 2\mathbb{1}_d \otimes \mathbb{1}_d - \sum_{\alpha=0}^d \sum_{i=0}^{d-1} |i_\alpha\rangle\langle i_\alpha| \otimes |i_\alpha\rangle\langle i_\alpha|. \tag{5}$$

We can then apply the 2-design property of a complete set of MUBs [78], i.e.

$$\sum_{\alpha=0}^d \sum_{i=0}^{d-1} |i_\alpha\rangle\langle i_\alpha| \otimes |i_\alpha\rangle\langle i_\alpha| = 2 \Pi_{\text{sym}}, \tag{6}$$

where $\Pi_{\text{sym}} = \frac{1}{2}(\mathbb{1}_d \otimes \mathbb{1}_d + \mathbb{F}_d)$ denotes a projector onto the symmetric subspace of $\mathcal{H} \otimes \mathcal{H}$. Thus, we have $\mathbf{W}(\mathcal{M}_{d+1}, 0) = 2(\mathbb{1}_d \otimes \mathbb{1}_d - \Pi_{\text{sym}})$. Here, \mathbb{F}_d denotes a flip (swap) operator on $\mathbb{C}^d \otimes \mathbb{C}^d$ defined by

$$\mathbb{F}_d = \sum_{i,j=0}^{d-1} |i\rangle\langle j| \otimes |j\rangle\langle i|. \tag{7}$$

Actually, \mathbb{F}_d does not depend on the basis. One has, therefore,

$$\mathbf{W}(\mathcal{M}_{d+1}, 0) = 2 \Pi_{\text{asym}}, \tag{8}$$

where $\Pi_{\text{asym}} = \frac{1}{2}(\mathbb{1}_d \otimes \mathbb{1}_d - \mathbb{F}_d)$ denotes a projector onto the antisymmetric subspace in $\mathcal{H} \otimes \mathcal{H}$. Clearly, $\mathbf{W}(\mathcal{M}_{d+1}, 0) \geq 0$, and it is therefore not an EW (recall, that a witness \mathbf{W} has to

satisfy $\text{tr}[\mathbf{W}\rho_{\text{sep}}] \geq 0$ and must possess at least one strictly negative eigenvalue). However, its partial transpose

$$\mathbf{W}^\Gamma(\mathcal{M}_{d+1}, 0) = \mathbb{1}_d \otimes \mathbb{1}_d - dP_d^+, \tag{9}$$

where

$$P_d^+ = \frac{1}{d} \mathbb{F}_d^\Gamma = \frac{1}{d} \sum_{i,j=0}^{d-1} |i\rangle\langle j| \otimes |i\rangle\langle j|, \tag{10}$$

defines a decomposable EW corresponding to the well known reduction map [43, 79, 80]:

$$R(X) = \mathbb{1}_d \text{Tr}X - X. \tag{11}$$

It proves that for a trivial permutation, i.e. $s = 0$, and access to a complete set of $d + 1$ MUBs, one cannot detect bound entangled states. Instead, it detects, e.g. all possible entangled states of the family of isotropic states $\rho_{\text{iso}} = \frac{1-p}{d^2} \mathbb{1}_{d^2} + p P_{k\ell}$ (where $P_{k\ell}$ are the maximally entangled Bell states, defined in equation (19)).

4. Full set of $d + 1$ MUBs and $s > 0$

To construct non-decomposable witnesses one needs to consider a nontrivial shift $s > 0$ (s is defined mod d), i.e.

$$\mathbf{W}(\mathcal{M}_{d+1}, s) = 2\mathbb{1}_d \otimes \mathbb{1}_d - \mathbf{B}(\mathcal{M}_{d+1}, s). \tag{12}$$

Using the two-design property one immediately finds

$$\begin{aligned} \mathbf{W}(\mathcal{M}_{d+1}, s) &= \mathbb{1}_d \otimes \mathbb{1}_d + (\Pi_0 - \Pi_s) - \mathbb{F}_d \\ &= 2\Pi_{\text{asym}} + (\Pi_0 - \Pi_s), \end{aligned} \tag{13}$$

where

$$\Pi_s = \sum_{i=0}^{d-1} |i\rangle\langle i| \otimes |i+s\rangle\langle i+s|. \tag{14}$$

Note, that $\Pi_k \Pi_\ell = \delta_{k\ell} \Pi_k$, and $\Pi_0 + \dots + \Pi_{d-1} = \mathbb{1}_d \otimes \mathbb{1}_d$. Interestingly, one has

$$\mathbf{W}(\mathcal{M}_{d+1}, s) = \mathbf{W}(\mathcal{M}_{d+1}, 0) + \Pi_0 - \Pi_s, \tag{15}$$

or, equivalently

$$\mathbf{W}(\mathcal{M}_{d+1}, s) + \Pi_s = \mathbf{W}(\mathcal{M}_{d+1}, 0) + \Pi_0. \tag{16}$$

Now, we investigate what happens if we consider the transpose. Indeed, it turns out that $\mathbf{W}^\Gamma(\mathcal{M}_{d+1}, s)$ is a Bell diagonal bipartite operator. To show this, let us introduce a set of unitary Weyl operators

$$U_{k\ell}|m\rangle = \omega^{k(m-\ell)}|m-\ell\rangle, \tag{17}$$

where $\omega = e^{2\pi i/d}$. Note that $U_{00} = \mathbb{1}_d$, $\text{tr}[U_{k\ell}U_{mn}^\dagger] = d \delta_{km} \delta_{\ell n}$, and

$$U_{mn}U_{k\ell} = \omega^{nk}U_{m+k, n+\ell}, \quad U_{k\ell}^\dagger = \omega^{k\ell}U_{-k, -\ell}. \tag{18}$$

We define the rank-1 projectors, corresponding to maximally entangled states,

$$P_{k\ell} := |\psi_{k\ell}\rangle\langle\psi_{k\ell}|, \tag{19}$$

where

$$|\psi_{k\ell}\rangle = \mathbb{1}_d \otimes U_{k\ell} |\psi_d^+\rangle, \tag{20}$$

and $|\psi_d^+\rangle = \frac{1}{\sqrt{d}} \sum_{n=0}^{d-1} |n\rangle \otimes |n\rangle$ denotes a canonical maximally entangled state. A set of d^2 vectors $|\psi_{k\ell}\rangle$ —called generalised Bell vectors—defines an orthonormal basis in $\mathcal{H} \otimes \mathcal{H}$. Note that $\Pi_\ell := P_{0\ell} + P_{1\ell} + \dots + P_{d-1,\ell}$, and whence

$$\begin{aligned} \mathbf{W}^\Gamma(\mathcal{M}_{d+1}, s) &= \mathbb{1}_d \otimes \mathbb{1}_d + (\Pi_0 - \Pi_s) - dP_d^+ \\ &= \mathbb{1}_d \otimes \mathbb{1}_d + \sum_{k=0}^{d-1} (P_{k0} - P_{ks}) - dP_d^+ \\ &= \mathbf{W}_{\text{Bell}}(s). \end{aligned} \tag{21}$$

Now we show that the operator $\mathbf{W}_{\text{Bell}}(s)$ defines a witness capable of detecting bound entanglement if the shift is nontrivial. In particular:

Proposition 1. *Let $s > 0$ and $d > 2$. Then $\mathbf{W}_{\text{Bell}}(s)$ defines a non-decomposable EW if and only if $s \neq d/2$.*

Proof. Consider the following state,

$$\rho_x = \frac{1}{\mathcal{N}} \left([\mathbb{1}_d \otimes \mathbb{1}_d - \Pi_0 - \Pi_s - \Pi_{d-s}] + \frac{1}{x} \Pi_s + x \Pi_{d-s} + dP_{00} \right), \tag{22}$$

with \mathcal{N} being the normalisation factor, and $x > 0$. This construction requires $s \neq d - s$, that is, $d \neq 2s$. Moreover, $s \rightarrow d - s$ corresponds to $x \rightarrow \frac{1}{x}$. Note, that ρ_x is a PPT state. Indeed, positivity of ρ_x follows from

$$\mathbb{1}_d \otimes \mathbb{1}_d - \Pi_0 - \Pi_s - \Pi_{d-s} \geq 0,$$

since Π_k are mutually orthogonal projectors. Now, to show that $\rho_x^\Gamma \geq 0$, let us observe due to $\Pi_k^\Gamma = \Pi_k$ (these are diagonal matrices), one has

$$\rho_x^\Gamma = \frac{1}{\mathcal{N}} \left([\mathbb{1}_d \otimes \mathbb{1}_d - \Pi_0 - \Pi_s - \Pi_{d-s}] + \frac{1}{x} \Pi_s + x \Pi_{d-s} + dP_{00}^\Gamma \right).$$

Recalling, that $dP_{00}^\Gamma = \mathbb{F}_d = \sum_{k,l} |k\rangle\langle l| \otimes |l\rangle\langle k|$, we observe that ρ_x^Γ consists of 2×2 blocks corresponding to the following 2×2 matrices

$$\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}, \quad \begin{pmatrix} x & 1 \\ 1 & 1/x \end{pmatrix}.$$

Clearly, both matrices are positive definite and, hence, the positivity of ρ_x^Γ follows.

Straightforwardly, we obtain

$$\text{tr}[\mathbf{W}_{\text{Bell}}(s)\rho_x] = \frac{1}{\mathcal{N}}(d(d-2) + dx - d(d-1)) = \frac{d}{\mathcal{N}}(x-1), \tag{23}$$

and, hence, $\text{tr}[\mathbf{W}^\Gamma(\mathcal{M}_{d+1}, s)\rho_x] < 0$ whenever $x < 1$. Thus, $\mathbf{W}_{\text{Bell}}(s)$ detects the PPT entangled state ρ_x with $x < 1$ and $\mathbf{W}_{\text{Bell}}(s)$ is a non-decomposable EW.

Now, if $s = \frac{d}{2}$, then $\mathbf{W}_{\text{Bell}}(s) = A(s) + B^\Gamma(s)$, where

$$A(s) = \sum_{n=0}^{s-1} (|nn\rangle - |(n+s)(n+s)\rangle)(\langle nn| - \langle (n+s)(n+s)|), \tag{24}$$

and

$$B(s) = \sum_{i < j \neq i+s} (|ij\rangle - |ji\rangle)(\langle ij| - \langle ji|). \tag{25}$$

Clearly, $A(s) \geq 0$ and $B(s) \geq 0$ (being sums of rank-1 projectors) which proves that $\mathbf{W}_{\text{Bell}}(s)$ is decomposable. \square

5. $d + 1$ MUBs are not necessary to detect bound entanglement

Interestingly, the full set of $d + 1$ MUBs is not necessary to detect bound entanglement. The main result of our paper consists of the following:

Theorem 1. *Let $2s \neq d$, with $s > 0$ and $d > 2$. If $m > \frac{d}{2} + 1$, then $\mathbf{W}^\Gamma(\mathcal{M}_m, s)$ defines a non-decomposable EW, i.e. detects bound entangled states.*

Proof. Let us start by writing the witness as

$$\mathbf{W}^\Gamma(\mathcal{M}_m, s) = \sum_{i,j,k,l=0}^{d-1} W_{ij;kl} |i\rangle\langle j| \otimes |l\rangle\langle k|, \tag{26}$$

with $W_{ij;kl}$ the elements of a matrix W . It is clear that the matrix elements $W_{ij;kl}$ depend on a particular set \mathcal{M}_m of MUBs. One can, however, find a subset of $\{W_{ij;kl}\}$ which is universal, i.e. independent of the choice of MUBs. In particular, the subset of elements

$$W_{ii;jj} = \begin{cases} 1 & \text{for } j \neq i + s, \\ 0 & \text{for } j = i + s, \end{cases} \tag{27}$$

and

$$W_{ij;ij} = -\frac{m-1}{d}, \text{ for } i \neq j, \tag{28}$$

is universal for any set \mathcal{M}_m of m MUBs in \mathbb{C}^d . This is a consequence of the property,

$$\text{tr} \left[\sum_{i=0}^{d-1} (|i_\alpha\rangle\langle i_\alpha| \otimes |i_\alpha^*\rangle\langle i_\alpha^*|) (|\ell\rangle\langle k| \otimes |\ell\rangle\langle k|) \right] = \frac{1}{d}, \tag{29}$$

which holds for any pair k and ℓ , and a fixed number of m MUBs.

Suppose that $\mathbf{W}^\Gamma(\mathcal{M}_m, s)$ is decomposable. That is, for some $A, B \geq 0$ we have $W = A + B^\Gamma$. Consider three 2×2 submatrices of W from (26):

$$\begin{aligned} A_1 &= \begin{pmatrix} W_{00;ss} & W_{0s;0s} \\ W_{s0;s0} & W_{ss;00} \end{pmatrix} = \begin{pmatrix} 0 & a \\ a & 1 \end{pmatrix} \\ A_2 &= \begin{pmatrix} W_{00;rr} & W_{0r;0r} \\ W_{r0;r0} & W_{rr;00} \end{pmatrix} = \begin{pmatrix} 1 & a \\ a & 0 \end{pmatrix} \\ A_3 &= \begin{pmatrix} W_{ss;rr} & W_{sr;sr} \\ W_{rs;rs} & W_{rr;ss} \end{pmatrix} = \begin{pmatrix} 1 & a \\ a & 0 \end{pmatrix} \end{aligned}$$

with $a = -(m-1)/d$ and $r = d - s$. Note that A_1, A_2 and A_3 are not positive definite. Therefore, if $\mathbf{W}^\Gamma(\mathcal{M}_m, s)$ is decomposable, then A_1, A_2 and A_3 must originate from B^Γ , and not from A which is positive definite. We therefore need to check whether B is positive, given that A_1, A_2 and A_3 are submatrices of B^Γ .

If one takes the partial transpose of B^Γ , the off diagonal elements of A_1, A_2 and A_3 that appear in B^Γ now form a 3×3 submatrix of B ,

$$B_{3 \times 3} = \begin{pmatrix} W_{00;00} & W_{0s;0s} & W_{0r;0r} \\ W_{s0;s0} & W_{ss;ss} & W_{sr;sr} \\ W_{r0;r0} & W_{rs;rs} & W_{rr;rr} \end{pmatrix} = \begin{pmatrix} 1 & a & a \\ a & 1 & a \\ a & a & 1 \end{pmatrix}. \tag{30}$$

It holds that $B_{3 \times 3} \geq 0$ if and only if $a \in [-\frac{1}{2}, 1]$. Recall that $a = -(m-1)/d$, so that we obtain a necessary condition for the decomposability, i.e. $m \leq \frac{d}{2} + 1$. This shows that, whenever

$$m > \frac{d}{2} + 1, \tag{31}$$

the witness $\mathbf{W}^\Gamma(\mathcal{M}_m, s)$ is not decomposable. □

Interestingly, the proof does not rely on the particular choice of m MUBs. Thus, any set of MUBs (that is sufficiently large) results in a non-decomposable witness able to detect bound entangled states.

6. Two MUBs give rise to a decomposable witness

We now show that a witness $\mathbf{W}^\Gamma(\mathcal{M}_2, s)$ consisting of two Fourier connected MUBs in \mathbb{C}^d gives rise to a decomposable witness. Suppose that \mathcal{M}_2 consists of the standard basis $\mathcal{B}_0 = \{|0\rangle, |1\rangle, \dots, |d-1\rangle\}$ and its Fourier transform $\mathcal{B}_1 = \{|\tilde{0}\rangle, |\tilde{1}\rangle, \dots, |\tilde{d-1}\rangle\}$, where

$$|\tilde{k}\rangle = \frac{1}{\sqrt{d}} \sum_{j=0}^{d-1} \omega^{kj} |j\rangle, \tag{32}$$

and $\omega = e^{2\pi i/d}$. Then, we define

$$\mathbf{W}^\Gamma(\mathcal{M}_2, s) = \frac{d+1}{d} \mathbb{1}_d \otimes \mathbb{1}_d - \sum_{\ell=0}^{d-1} |\ell\rangle\langle\ell| \otimes |\ell+s\rangle\langle\ell+s| - \sum_{k=0}^{d-1} |\tilde{k}\rangle\langle\tilde{k}| \otimes |\tilde{k}^*\rangle\langle\tilde{k}^*|. \tag{33}$$

One finds

$$\sum_{k=0}^{d-1} |\tilde{k}\rangle\langle\tilde{k}| \otimes |\tilde{k}^*\rangle\langle\tilde{k}^*| = \frac{1}{d} \sum_{n=0}^{d-1} \sum_{i,j=0}^{d-1} |i\rangle\langle i+n| \otimes |j\rangle\langle j+n|, \tag{34}$$

and hence

$$\mathbf{W}^\Gamma(\mathcal{M}_2, s) = \frac{1}{d}(A(s) + B^\Gamma(s)), \tag{35}$$

where

$$A(s) = (d-1)(\mathbb{1}_d \otimes \mathbb{1}_d - \Pi_s) - \sum_{\substack{n=0 \\ n \neq s}}^{d-1} \sum_{i \neq j=0}^{d-1} |i\rangle\langle j| \otimes |i+n\rangle\langle j+n|, \tag{36}$$

and $B(s)$ is a sum of $d(d-1)/2$ rank-1 projectors:

$$B(s) = \sum_{i \neq j=0}^{d-1} |\Psi_{ij}(s)\rangle\langle\Psi_{ij}(s)|, \tag{37}$$

where

$$|\Psi_{ij}(s)\rangle = |i, j + s\rangle - |j, i + s\rangle. \tag{38}$$

Evidently $A(s), B(s) \geq 0$, and hence $\mathbf{W}^\Gamma(\mathcal{M}_2, s)$ is decomposable for any s .

The proof also works for general pairs of Heisenberg-Weyl MUBs in prime dimensions $d = p$, however it is slightly more technical. For example, see the witnesses $\mathbf{W}_{(0,1)}$ and $\mathbf{W}_{(0,3)}$ defined in section 7. One may expect that a decomposition of the witness exists for *any* pair of MUBs, but a proof is hindered by the fact that the classification of pairs is an open problem. Hence, we are unable to describe the general structure of the witness and determine its decomposition. In fact, the classification of pairs of MUBs in \mathbb{C}^d is equivalent to a classification all $d \times d$ complex Hadamard matrices, which is currently known for $d \leq 5$ only [81].

The situation is further complicated by the fact that *equivalent* sets of MUBs (up to some unitary transformation) can give rise to different witnesses. For example, a unitary transformation connecting one set of MUBs to another may cause a permutation of the elements of the standard basis. Such a transformation would mean the witness no longer has the particular form given in equation (4), and the resulting effect on its (non-)decomposability is unclear.

7. Example: $d = 3$

If $d = 3$, condition (31) implies that three MUBs are sufficient to detect bound entanglement. Actually, we can also show that this condition is necessary. Consider the complete set of four Heisenberg-Weyl MUBs, $\mathcal{B}_0 = \{|0\rangle, |1\rangle, |2\rangle\}$ together with $\mathcal{B}_1, \mathcal{B}_2$ and \mathcal{B}_3 , which we represent by the following matrices,

$$B_1 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^* \\ 1 & \omega^* & \omega \end{pmatrix}, B_2 = \frac{1}{\sqrt{3}} \begin{pmatrix} 1 & 1 & 1 \\ 1 & \omega & \omega^* \\ \omega^* & \omega & 1 \end{pmatrix},$$

and $B_3 = B_2^*$, where the columns of B_1, B_2 and B_3 correspond to the basis vectors of $\mathcal{B}_1, \mathcal{B}_2$ and \mathcal{B}_3 , respectively. One finds three EWs,

$$\mathbf{W}^\Gamma(\{\mathcal{B}_i, \mathcal{B}_j, \mathcal{B}_k\}, 1) := \mathbf{W}_{(i,j,k)},$$

given by

$$\mathbf{W}_{(0,1,2)} = \left(\begin{array}{ccc|ccc|ccc} 1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & 1 & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \hline \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \hline \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1 & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1 \end{array} \right),$$

$$\mathbf{W}_{(0,1,3)} = \mathbf{W}_{(0,1,2)}^*,$$

and

$$\mathbf{W}_{(0,2,3)} = \left(\begin{array}{ccc|ccc|ccc} 1 & \cdot & \cdot & \cdot & -\frac{2}{3} & \cdot & \cdot & \cdot & -\frac{2}{3} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \frac{1}{3} & \cdot & \cdot \\ \cdot & \cdot & 1 & \frac{1}{3} & \cdot & \cdot & \cdot & \frac{1}{3} & \cdot \\ \hline \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot & \frac{1}{3} & \cdot \\ -\frac{2}{3} & \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot & -\frac{2}{3} \\ \cdot & \frac{1}{3} & \cdot & \cdot & \cdot & \cdot & \frac{1}{3} & \cdot & \cdot \\ \hline \cdot & \frac{1}{3} & \cdot & \cdot & \cdot & \frac{1}{3} & \cdot & \cdot & \cdot \\ \cdot & \cdot & \frac{1}{3} & \frac{1}{3} & \cdot & \cdot & \cdot & 1 & \cdot \\ -\frac{2}{3} & \cdot & \cdot & \cdot & -\frac{2}{3} & \cdot & \cdot & \cdot & 1 \end{array} \right),$$

where the dots represent zeros.

It is, therefore, clear that the matrix structure of $\mathbf{W}^\Gamma(\mathcal{M}_3, 1)$ depends on a particular set of \mathcal{M}_3 . However, all three witnesses are non-decomposable and they detect the same PPT entangled state,

$$\rho_x = \frac{1}{\mathcal{N}} \left(\begin{array}{ccc|ccc|ccc} 1 & \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot & 1 \\ \cdot & 1/x & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \cdot & \cdot & x & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \hline \cdot & \cdot & \cdot & x & \cdot & \cdot & \cdot & \cdot & \cdot \\ 1 & \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot & 1 \\ \cdot & \cdot & \cdot & \cdot & \cdot & 1/x & \cdot & \cdot & \cdot \\ \hline \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & 1/x & \cdot & \cdot \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & x & \cdot \\ 1 & \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot & 1 \end{array} \right), \tag{39}$$

with $x \neq 1$ (for $x = 1$ it is separable).

Similarly, one finds three witnesses $\mathbf{W}^\Gamma(\{\mathcal{B}_i, \mathcal{B}_j\}, 1) := \mathbf{W}_{(i,j)}$ as follows:

$$\mathbf{W}_{(0,1)} = \left(\begin{array}{ccc|ccc|ccc} 1 & \cdot & \cdot & \cdot & -\frac{1}{3} & \cdot & \cdot & \cdot & -\frac{1}{3} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & -\frac{1}{3} & \cdot & \cdot \\ \cdot & \cdot & 1 & -\frac{1}{3} & \cdot & \cdot & \cdot & -\frac{1}{3} & \cdot \\ \hline \cdot & \cdot & -\frac{1}{3} & 1 & \cdot & \cdot & \cdot & -\frac{1}{3} & \cdot \\ -\frac{1}{3} & \cdot & \cdot & \cdot & 1 & \cdot & \cdot & \cdot & -\frac{1}{3} \\ \cdot & -\frac{1}{3} & \cdot & \cdot & \cdot & \cdot & \cdot & -\frac{1}{3} & \cdot \\ \hline \cdot & -\frac{1}{3} & \cdot & \cdot & -\frac{1}{3} & \cdot & -\frac{1}{3} & \cdot & \cdot \\ \cdot & \cdot & -\frac{1}{3} & -\frac{1}{3} & \cdot & \cdot & \cdot & 1 & \cdot \\ -\frac{1}{3} & \cdot & \cdot & \cdot & -\frac{1}{3} & \cdot & \cdot & \cdot & 1 \end{array} \right)$$

$$\mathbf{W}_{(0,3)} = \left(\begin{array}{ccc|ccc|ccc} -1 & \cdot & \cdot & \cdot & \frac{1}{3} & \cdot & \cdot & \cdot & \frac{1}{3} \\ \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \frac{1}{3} & \cdot & \cdot \\ \cdot & \cdot & -1 & \frac{\mathcal{E}^*}{3} & \cdot & \cdot & \cdot & \frac{\mathcal{E}^*}{3} & \cdot \\ \hline \cdot & \cdot & \frac{\mathcal{E}}{3} & -1 & \cdot & \cdot & \cdot & \frac{\mathcal{E}}{3} & \cdot \\ \frac{1}{3} & \cdot & \cdot & \cdot & -1 & \cdot & \cdot & \cdot & \frac{1}{3} \\ \cdot & \frac{\mathcal{E}^*}{3} & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot & \cdot \\ \hline \cdot & \frac{\mathcal{E}}{3} & \cdot & \cdot & \cdot & \cdot & \frac{\mathcal{E}^*}{3} & \cdot & \cdot \\ \cdot & \cdot & \frac{\mathcal{E}^*}{3} & \cdot & \cdot & \cdot & \cdot & -1 & \cdot \\ \frac{1}{3} & \cdot & \cdot & \cdot & \frac{1}{3} & \cdot & \cdot & \cdot & -1 \end{array} \right)$$

and

$$\mathbf{W}_{(0,2)} = \mathbf{W}_{(0,3)}^*.$$

Again, the matrix structure of $\mathbf{W}(\mathcal{M}_2, 1)$ depends on the particular set of \mathcal{M}_2 . However, all three witnesses are decomposable. In particular $\mathbf{W}_{(0,1)} = A + B^\Gamma$, where

$$A = \frac{1}{3} \left(\begin{array}{ccc|ccc} 2 & . & . & . & -1 & . & . & . & -1 \\ . & . & . & . & . & . & . & . & . \\ . & . & 2 & -1 & . & . & . & -1 & . \\ \hline . & . & -1 & 2 & . & . & . & -1 & . \\ -1 & . & . & . & 2 & . & . & . & -1 \\ . & . & . & . & . & . & . & . & . \\ \hline . & . & . & . & . & . & . & . & . \\ . & . & -1 & -1 & . & . & . & 2 & . \\ -1 & . & . & . & -1 & . & . & . & 2 \end{array} \right) \quad (40)$$

and

$$B = \frac{1}{3} \left(\begin{array}{ccc|ccc} 1 & . & . & . & . & . & . & -1 & . \\ . & . & . & . & . & . & . & . & . \\ . & . & 1 & . & -1 & . & . & . & . \\ \hline . & . & . & 1 & . & . & . & . & -1 \\ . & . & -1 & . & 1 & . & . & . & . \\ . & . & . & . & . & . & . & . & . \\ \hline . & . & . & . & . & . & . & . & . \\ -1 & . & . & . & . & . & . & 1 & . \\ . & . & . & -1 & . & . & . & . & 1 \end{array} \right), \quad (41)$$

with $A, B \geq 0$. Similar decompositions hold for the remaining two witnesses.

8. Special dimensions: $d = 2^r$

We now improve Theorem 1 by showing that 3 MUBs (i.e. $m = \frac{d}{2} + 1$) are sufficient for bound entanglement detection when $d = 4$. In particular, we find specific examples of three MUBs which detect bound entanglement. It remains open whether all triples (for which there is an infinite family [82]) are sufficient for bound entanglement detection.

Consider the following set of three MUBs in \mathbb{C}^4 , denoted by $\{\mathcal{B}_0, \mathcal{B}_1, \mathcal{B}_x\}$, where \mathcal{B}_0 is the canonical basis, \mathcal{B}_1 , in matrix form, is given by

$$B_1 = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & -1 & 1 \\ 1 & -1 & 1 & -1 \end{pmatrix}, \quad (42)$$

and \mathcal{B}_x , a third mutually unbiased basis, is chosen as either

$$B_{\text{ext}} = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ i & -i & i & -i \\ -1 & -1 & 1 & 1 \\ i & -i & -i & i \end{pmatrix}, \quad (43)$$

or

$$B_{\text{unext}} = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ -1 & 1 & 1 & -1 \\ 1 & -1 & 1 & -1 \end{pmatrix}. \quad (44)$$

The three MUBs $\{\mathcal{B}_0, \mathcal{B}_1, \mathcal{B}_{\text{ext}}\}$ form an *extendible* set, which is a subset of the complete set of five Heisenberg-Weyl MUBs. In contrast, $\{\mathcal{B}_0, \mathcal{B}_1, \mathcal{B}_{\text{unext}}\}$ is *unextendible*, in the sense that no other basis is mutually unbiased to all three bases [82]. We label the associated EWs

$$\mathbf{W}^\Gamma(\{\mathcal{B}_0, \mathcal{B}_1, \mathcal{B}_x\}, 1) := \mathbf{W}_x. \quad (45)$$

Now we have the following result.

Proposition 2. *The EWs*

$$\mathbf{W}_{\text{ext}} \text{ and } \mathbf{W}_{\text{unext}}$$

are non-decomposable.

Proof. In the [appendix](#) (equations (46) and (48)) we present the matrix form of these witnesses. Non-decomposability follows by constructing families of PPT states ρ_a and ρ_b , defined in (47) and (49), which violate the separability criterion. Firstly, we see that

$$\text{tr}[\mathbf{W}_{\text{ext}}\rho_a] = \frac{4}{\mathcal{N}}(a - 1) < 0,$$

for $a < 1$. Since ρ_a is PPT in this region and the separability condition is clearly violated when $a < 1$, this proves that \mathbf{W}_{ext} is non-decomposable.

Similarly, after a simple calculation, we have

$$\text{tr}[\mathbf{W}_{\text{unext}}\rho_b] = \frac{4}{\mathcal{N}}(b - 1) < 0,$$

for $b < 1$. This proves that $\mathbf{W}_{\text{unext}}$ is indeed non-decomposable (and ρ_b is PPT entangled for $b < 1$).

Hence, for $d = 4$ we have shown that both extendible and unextendible MUBs give rise to a non-decomposable EWs. As a by-product we present two new families (47) and (49) of PPT entangled states in a 4×4 dimensional system that are not locally unitarily equivalent. \square

Furthermore, from section 6, the EW $\mathbf{W}^\Gamma(\mathcal{M}_2, 1)$, with a Fourier connected pair of measurements, is decomposable. Hence, while we have not ruled out all possible pairs from detecting bound entanglement, the extension to a third basis appears necessary. When $d = 2^r$, we predict that $\frac{d}{2} + 1$ MUBs are both necessary and sufficient for the witness $\mathbf{W}^\Gamma(\mathcal{M}_m, s)$ to be non-decomposable.

Conjecture 1. If $d = 2^r$ and $r > 1$, the minimal number of MUBs required for bound entanglement detection equals $m = \frac{d}{2} + 1 = 2^{r-1} + 1$.

We point out that the three MUBs used to construct $\mathbf{W}_{\text{unext}}$ have the unusual feature of being bases in \mathbb{R}^4 . It is known that the number of *real* MUBs in \mathbb{R}^d is at most $\frac{d}{2} + 1$, and the bound is tight when $d = 4^r$ [83]. This suggests the possibility of finding non-decomposable witnesses from real MUBs in these special dimensions. It would be interesting to determine the usefulness of real MUBs for entanglement detection [84], especially in light of their various constructions, e.g. [83].

The above conjecture, together with the example of the witness $\mathbf{W}_{\text{unext}}$, suggests that *large* sets of unextendible MUBs in higher dimensions can be used for bound entanglement detection. For $d = 4, 8$ it is known that $m = d/2 + 1$ unextendible MUBs exist, and it is conjectured that such bases also occur when $d = 2^r$ [85]. In contrast, smaller sets of unextendible MUBs are not expected to yield examples of non-decomposable witnesses, including those constructed up to $d = 16$ in [86], and the $p + 1$ unextendible MUBs found in [87] when $d = p^2$.

9. Conclusions

In this article we investigated the minimal number of MUBs needed to detect bound entangled states. We have shown that sets of m MUBs in \mathbb{C}^d , when $m > d/2 + 1$, give rise to non-decomposable witnesses. Thus, bound entanglement verification can be achieved with significantly fewer measurements than a complete set of $d + 1$ MUBs. This presents a more favourable, experimentally friendly, detection approach compared with typical quantum state tomography methods, or witnesses composed of complete sets. It also opens up the possibility of applying these witnesses to detect PPT entangled states in dimensions *without* the existence of a complete set, i.e. in non-prime-power dimensions.

We have shown that the original EW composed of $d + 1$ MUBs in [40] is decomposable, and therefore cannot detect bound entangled states. We also found that pairs of Fourier connected MUBs (as well as Heisenberg-Weyl pairs in prime dimensions) are unable to certify bound entanglement. When $d = 3$, we proved that three MUBs are necessary and sufficient to detect bound entangled states. In $d = 4$, we strengthened theorem 1 to show that PPT entangled states can be detected with $m = 3 = \frac{d}{2} + 1$ MUBs. As a consequence, we found new families of bipartite bound entangled states in a (4×4) system, and conjectured $m = \frac{d}{2} + 1$ MUBs are required to detect bound entanglement if $d = 2^r$ with $r > 1$.

In recent work it has been shown that certain sets of MUBs are more useful than others—a consequence of the unitary inequivalence of different classes of MUBs—in information processing tasks [42, 88, 89]. In contrast, theorem 1 shows that since *any* set of m MUBs ($m > d/2 + 1$) detects bound entangled states, the non-decomposability of a witness is a universal property of MUBs. However, since different equivalence classes give rise to different EWs, an interesting interplay is revealed between the universality of MUBs for non-decomposable witnesses and the more local dependence on the choice of MUBs in detecting a given bound entangled state. The nature of such connections would be worth investigating further since they may shed light on the structure of the set of bound entangled states.

Other interesting directions include studying the optimality of these witnesses, and determining whether a reduction in the number of measurements can result in a decomposable witness transitioning to a non-decomposable one. Further open questions include whether *every* pair of MUBs is unable to detect bound entanglement, and, more generally, the minimal number of MUBs needed to detect bound entangled states. Another avenue to explore is the construction

of new families of bipartite PPT entangles states, which are detected by the large class of non-decomposable EWs we have studied.

Data availability statement

All data that support the findings of this study are included within the article (and any supplementary files).

Acknowledgments

J B is supported by the National Research Foundation of Korea (Grant Nos. NRF-2021R1A2C200 6309, NRF-2022M1A3C2069728) and the Institute for Information & Communication Technology Promotion (IITP) (the ITRC Program/IITP-2022-2018-0-01402). A B and D C were supported by the Polish National Science Centre Project 2018/30/A/ST2/00837. B C H acknowledges gratefully that this research was funded in whole, or in part, by the Austrian Science Fund (FWF) Project P36102-N. D M has received funding from the European Union’s Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie Grant Agreement No. 663830. D M also acknowledges financial support by the TEAM-NET project co-financed by the E U within the Smart Growth Operational Programme (Contract No. POIR.04.04.00-00-17C1/18-00).

Appendix. Non-decomposable witness when $d = 4$

As shown in proposition 2, we find two non-decomposable EWs using sets of three MUBs. The witness corresponding to the three Heisenberg–Weyl MUBs is given by

$$W_{\text{ext}} = \begin{pmatrix}
 \begin{array}{cccc|cccc|cccc|cccc}
 1 & . & . & . & . & -\frac{1}{2} & . & . & . & . & . & . & -\frac{1}{2} & . & . & . & . \\
 . & . & . & . & . & . & . & . & . & . & . & . & . & . & -\frac{1}{2} & . & . \\
 . & . & 1 & . & . & . & . & . & . & -\frac{1}{2} & . & . & . & . & . & . & . \\
 . & . & . & 1 & . & . & -\frac{1}{2} & . & . & . & . & . & . & . & . & . & . \\
 \hline
 . & . & . & . & . & 1 & . & . & . & . & . & -\frac{1}{2} & . & . & . & . & . \\
 -\frac{1}{2} & . & . & . & . & . & 1 & . & . & . & . & -\frac{1}{2} & . & . & . & . & -\frac{1}{2} \\
 . & . & . & -\frac{1}{2} & . & . & . & . & . & . & . & . & . & . & . & . & . \\
 . & . & . & . & . & . & . & . & 1 & . & . & . & . & -\frac{1}{2} & . & . & . \\
 \hline
 . & . & -\frac{1}{2} & . & . & . & . & . & . & 1 & . & . & . & . & . & . & . \\
 . & . & . & . & . & . & . & . & . & . & 1 & . & . & -\frac{1}{2} & . & . & . \\
 -\frac{1}{2} & . & . & . & . & -\frac{1}{2} & . & . & . & . & . & 1 & . & . & . & -\frac{1}{2} & . \\
 . & . & . & . & -\frac{1}{2} & . & . & . & . & . & . & . & . & . & . & . & . \\
 \hline
 . & . & . & . & . & . & . & . & . & -\frac{1}{2} & . & . & . & . & . & . & . \\
 . & . & . & . & . & . & . & . & -\frac{1}{2} & . & . & . & . & 1 & . & . & . \\
 . & -\frac{1}{2} & . & . & . & . & . & . & . & . & . & . & . & . & 1 & . & . \\
 -\frac{1}{2} & . & . & . & . & -\frac{1}{2} & . & . & . & . & -\frac{1}{2} & . & . & . & . & . & 1
 \end{array}
 \end{pmatrix}. \tag{46}$$

- [19] Banerjee S, Alok A K, Srikanth R and Hiesmayr B C 2015 *Eur. Phys. J. C* **75** 1–9
- [20] Gurvits L 2003 *Proc. 35th Annual ACM Symp. on Theory of Computing* pp 10–19
- [21] Chruściński D and Pittenger A O 2008 *J. Phys. A: Math. Theor.* **41** 385301
- [22] Marconi C, Aloy A, Tura J and Sanpera A 2021 *Quantum* **5** 561
- [23] Horodecki P 1997 *Phys. Lett. A* **232** 333
- [24] Bennett C H, DiVincenzo D P, Mor T, Shor P W, Smolin J A and Terhal B M 1999 *Phys. Rev. Lett.* **82** 5385
- [25] DiVincenzo D P, Mor T, Shor P W, Smolin J A and Terhal B M 2003 *Commun. Math. Phys.* **238** 379
- [26] Bertlmann R A and Krammer P 2009 *Ann. Phys.* **324** 1388–407
- [27] Lockhart J, Gühne O and Severini S 2018 *Phys. Rev. A* **97** 062340
- [28] Bruß D and Peres A 2000 *Phys. Rev. A* **61** 030301
- [29] Hiesmayr B C and Löffler W 2014 *Phys. Scr.* **T160** 014017
- [30] Tura J, Aloy A, Quesada R, Lewenstein M and Sanpera A 2018 *Quantum* **2** 45
- [31] Šupić I and Bowles J 2020 *Quantum* **4** 337
- [32] Kliesch M and Roth I 2021 *PRX Quantum* **2** 010201
- [33] Bowles J, Šupić I, Cavalcanti D and Acín A 2018 *Phys. Rev. Lett.* **121** 180503
- [34] Branciard C, Rosset D, Liang Y-C and Gisin N 2013 *Phys. Rev. Lett.* **110** 060405
- [35] Verbanis E, Martin A, Rosset D, Lim C C W, Thew R T and Zbinden H 2016 *Phys. Rev. Lett.* **116** 190501
- [36] Durt T, Englert B-G, Bengtsson I and Życzkowski K 2010 *Int. J. Quantum Inf.* **8** 535
- [37] Ivanović I D 1981 *J. Phys. A* **14** 3241
- [38] Wootters W K and Fields B D 1989 *Ann. Phys., NY* **191** 363
- [39] Boykin P O, Sitharam M, Tiep P H and Wocjan P 2007 *Quantum Inf. Comput.* **7** 371–82
- [40] Spengler C, Huber M, Brierley S, Adaktylos T and Hiesmayr B C 2012 *Phys. Rev. A* **86** 022311
- [41] Bae J, Hiesmayr B C and McNulty D 2019 *New J. Phys.* **21** 013012
- [42] Hiesmayr B C, McNulty D, Baek S, Singha Roy S, Bae J and Chruściński D 2021 *New J. Phys.* **23** 093018
- [43] Chruściński D, Sarbicki G and Wudarski F 2018 *Phys. Rev. A* **97** 032318
- [44] Kalev A and Bae J 2013 *Phys. Rev. A* **87** 062314
- [45] Erker P, Krenn M and Huber M 2017 *Quantum* **1** 22
- [46] Kaniewski J, Šupić I, Tura J, Baccari F, Salavrakos A and Augusiak R 2019 *Quantum* **3** 198
- [47] Kalev A and Gour G 2014 *New J. Phys.* **16** 053038
- [48] Bavaresco J, Valencia N H Klöckl C, Pivoluska M, Erker P, Friis N, Malik M and Huber M 2018 *Nat. Phys.* **14** 1032
- [49] Hiesmayr B C and Löffler W 2013 *New J. Phys.* **15** 083036
- [50] Terhal B M 2001 *Linear Algebr. Appl.* **323** 61
- [51] Lewenstein M, Kraus B, Cirac J and Horodecki P 2000 *Phys. Rev. A* **62** 052310
- [52] Lewenstein M, Kraus B, Horodecki P and Cirac J I 2001 *Phys. Rev. A* **63** 044304
- [53] Ha K-C and Kye S-H 2012 *Phys. Rev. A* **86** 034301
- [54] Amselem E and Bourennane M 2009 *Nat. Phys.* **5** 748
- [55] Lavoie J, Kaltenbaek R, Piani M and Resch K J 2010 *Phys. Rev. Lett.* **105** 130501
- [56] Barreiro J T, Schindler P, Gühne O, Monz T, Chwalla M, Roos C F, Hennrich M and Blatt R 2010 *Nat. Phys.* **6** 943
- [57] Kampermann H, Bruß D, Peng X and Suter D 2010 *Phys. Rev. A* **81** 040304
- [58] Kaneda F, Shimizu R, Ishizaka S, Mitsumori Y, Kosaka H and Edamatsu K 2012 *Phys. Rev. Lett.* **109** 040501
- [59] DiGiuglielmo J, Sambrowski A, Hage B, Pineda C, Eisert J and Schnabel R 2011 *Phys. Rev. Lett.* **107** 240503
- [60] Giovannini D, Romero J, Leach J, Dudley A, Forbes A and Padgett M J 2013 *Phys. Rev. Lett.* **110** 143601
- [61] D’ambrosio V, Cardano F, Karimi E, Nagali E, Santamato E, Marrucci L and Sciarrino F 2013 *Sci. Rep.* **3** 2726
- [62] Bartkiewicz K, Černoč A, Lemr K, Miranowicz A and Nori F 2016 *Phys. Rev. A* **93** 062345
- [63] Mafu M, Dudley A, Goyal S, Giovannini D, McLaren M, Padgett M J, Konrad T, Petruccione F, Lütkenhaus N and Forbes A 2013 *Phys. Rev. A* **88** 032305
- [64] Adamson R B A and Steinberg A M 2010 *Phys. Rev. Lett.* **105** 030406
- [65] Lukens J M, Islam N T, Lim C C W and Gauthier D J 2018 *Appl. Phys. Lett.* **112** 111102
- [66] Hu X-M et al 2020 *Phys. Rev. Lett.* **125** 230501

- [67] Ecker S *et al* 2019 *Phys. Rev. X* **9** 041042
- [68] Herrera V N, Srivastav V, Pivoluska M, Huber M, Friis N, McCutcheon W and Malik M 2020 *Quantum* **4** 376
- [69] Kewming M J, Shrapnel S, White A G and Romero J 2020 *Phys. Rev. Lett.* **124** 250401
- [70] Bouchard F, Heshami K, England D, Fickler R, Boyd R W, Englert B-G, Sánchez-Soto L L and Karimi E 2018 *Quantum* **2** 111
- [71] Sentís G, Greiner J N, Shang J, Siewert J and Kleinmann M 2018 *Quantum* **2** 113
- [72] Schwemmer C, Knips L, Richart D, Weinfurter H, Moroder T, Kleinmann M and Gühne O 2015 *Phys. Rev. Lett.* **114** 080403
- [73] Roy A and Scott A J 2007 *J. Math. Phys.* **48** 072110
- [74] Rudolph O 2003 *Phys. Rev. A* **67** 032312
- [75] Tavakoli A, Farkas M, Rosset D, Bancal J-D and Kaniewski J 2021 *Sci. Adv.* **7** eabc3847
- [76] Zhu H, Hayashi M and Chen L 2016 *Phys. Rev. Lett.* **116** 070403
- [77] Bae J, Chruściński D and Hiesmayr B C 2020 *npj Quantum Inf.* **6** 15
- [78] Klappenecker A and Rötteler M 2005 *Proc. 2005 IEEE Int. Symp. on Information Theory* pp 1740–4
- [79] Cerf N J, Adami C and Gingrich R M 1999 *Phys. Rev. A* **60** 898
- [80] Horodecki M and Horodecki P 1999 *Phys. Rev. A* **59** 4206
- [81] Haagerup U 1997 *Operator Algebras and Quantum Field Theory* (Cambridge: International Press)
- [82] Brierley S, Weigert S and Bengtsson I 2010 *Quantum Inf. Comput.* **10** 803
- [83] Boykin P O, Sitharam M, Tarifi M and Wocjan P 2005 arXiv:0502024
- [84] Renou M-O, Trillo D, Weilenmann M, Le T P, Tavakoli A, Gisin N, Acín A and Navascués M 2021 *Nature* **600** 625
- [85] Mandayam P, Bandyopadhyay S, Grassl M and Wootters W K 2014 *Quantum Inf. Comput.* **14** 823
- [86] Grassl M, McNulty D, Mista D L Jr and Paterek T 2017 *Phys. Rev. A* **95** 012118
- [87] Hegde V and Mandayam P 2015 arXiv:1508.05892
- [88] Aguilar E A, Borkala J J, Mironowicz P and Pawłowski M 2018 *Phys. Rev. Lett.* **121** 050501
- [89] Designolle S, Skrzypczyk P, Fröwis F and Brunner N 2019 *Phys. Rev. Lett.* **122** 050402