

Perfect Matchings in Claw-free Cubic Graphs

Sang-il Oum*

Department of Mathematical Sciences
KAIST, Daejeon, 305-701, Republic of Korea

sangil@kaist.edu

Submitted: Nov 9, 2009; Accepted: Mar 7, 2011; Published: Mar 24, 2011

Mathematics Subject Classification: 05C70

Abstract

Lovász and Plummer conjectured that there exists a fixed positive constant c such that every cubic n -vertex graph with no cutedge has at least 2^{cn} perfect matchings. Their conjecture has been verified for bipartite graphs by Voorhoeve and planar graphs by Chudnovsky and Seymour. We prove that every claw-free cubic n -vertex graph with no cutedge has more than $2^{n/12}$ perfect matchings, thus verifying the conjecture for claw-free graphs.

1 Introduction

A graph is *claw-free* if it has no induced subgraph isomorphic to $K_{1,3}$. A graph is *cubic* if every vertex has exactly three incident edges. A well-known classical theorem of Petersen [10] states that every cubic graph with no cutedge has a perfect matching. Sumner [11] and Las Vergnas [7] independently showed that every connected claw-free graph with even number of vertices has a perfect matching. Both theorems imply that every claw-free cubic graph with no cutedge has at least one perfect matching.

In 1970s, Lovász and Plummer conjectured that every cubic graph with no cutedge has exponentially many perfect matchings; see [8, Conjecture 8.1.8]. The best lower bound has been obtained by Esperet, Kardoš, and Král' [6]. They showed that the number of perfect matchings in a sufficiently large cubic graph with no cutedge always exceeds any fixed linear function in the number of vertices.

So far the conjecture is known to be true for bipartite graphs and planar graphs. For bipartite graphs, Voorhoeve [12] proved that every *bipartite* cubic n -vertex graph has at least $6(4/3)^{n/2-3}$ perfect matchings. Recently, Chudnovsky and Seymour [2] proved that every *planar* cubic n -vertex graph with no cutedge has at least $2^{n/655978752}$ perfect matchings.

*Supported by Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Minister of Education, Science and Technology (2010-001655) and TJ Park Junior Faculty Fellowship.




Figure 1: Claw-free cubic graphs with only 9 perfect matchings

We prove that every claw-free cubic n -vertex graph with no cutedge has more than $2^{n/12}$ perfect matchings. The graph should not have any cutedge; in Figure 1, we provide an example of a claw-free cubic graph with only 9 perfect matchings.

Our approach is to use the structure of 2-edge-connected claw-free cubic graphs. The *cycle space* $\mathcal{C}(H)$ of H is a collection of the edge-disjoint union of cycles of H . It is well known that $\mathcal{C}(H)$ forms a vector space over $GF(2)$ and

$$\dim \mathcal{C}(H) = |E(H)| - |V(H)| + 1$$

if H is connected, see Diestel [3]. Roughly speaking, almost all 2-edge-connected claw-free cubic graph G can be built from a 2-edge-connected cubic multigraph H by certain operations so that members of $\mathcal{C}(H)$ can be mapped injectively to 2-factors of G . We will have two cases to consider; either H is big or small. If H is big, then $\mathcal{C}(H)$ is big enough to prove that G has many 2-factors. If H is small, then we find a 2-factor of H using many of the specified edges of H so that when transforming this 2-factor of H to that of G , each of those edges of H has many ways to make 2-factors of G .

2 Structure of 2-edge-connected claw-free cubic graphs

Graphs in this paper have no parallel edges and no loops, and multigraphs can have parallel edges and loops. We assume that a loop is counted twice when measuring a degree of a vertex in a multigraph. Every 2-edge-connected cubic multigraph cannot have loops because if it has a loop, then it must have a cutedge.

We describe the structure of claw-free cubic graphs given by Palmer et al. [9]. A *triangle* of a graph is a set of three pairwise adjacent vertices. *Replacing a vertex v with a triangle* in cubic graph is to replace v with three vertices v_1, v_2, v_3 forming a triangle so that if e_1, e_2, e_3 are three edges incident with v , then e_1, e_2, e_3 will be incident with v_1, v_2, v_3 respectively.

Every vertex in a claw-free cubic graph is in 1, 2, or 3 triangles. If a vertex is in 3 triangles, then the component containing the vertex is isomorphic to K_4 . If a vertex is in exactly 2 triangles, then it is in an induced subgraph isomorphic to $K_4 \setminus e$ for some edge e of K_4 . Such an induced subgraph is called a *diamond*. It is clear that no two distinct diamonds intersect.

A *string of diamonds* is a maximal sequence D_1, D_2, \dots, D_k of diamonds in which, for each $i \in \{1, 2, \dots, k-1\}$, D_i has a vertex adjacent to a vertex in D_{i+1} . A string of diamonds has exactly two vertices of degree 2, which are called the *head* and the *tail* of the string. *Replacing an edge $e = uv$ with a string of diamonds* with the head x and the tail y is to remove e and add edges ux and vy .

A connected claw-free cubic graph in which every vertex is in a diamond is called a *ring of diamonds*. We require that a ring of diamonds contains at least 2 diamonds. It is now straightforward to describe the structure of 2-edge-connected claw-free cubic graphs as follows.

Proposition 1. *A graph G is 2-edge-connected claw-free cubic if and only if either*

- (i) G is isomorphic to K_4 ,
- (ii) G is a ring of diamonds, or
- (iii) G can be built from a 2-edge-connected cubic multigraph H by replacing some edges of H with strings of diamonds and replacing each vertex of H with a triangle.

Proof. Let us first prove the “if” direction. It is easy to see that G is 2-edge-connected cubic and has no loops or parallel edges. If G is built as in (iii), then clearly G has neither loops nor parallel edges, and every vertex of G is in a triangle and therefore G is claw-free. Note that since H is 2-edge-connected, H cannot have loops.

To prove the “only if” direction, let us assume that G is a 2-edge-connected claw-free cubic graph. We may assume that G is not isomorphic to K_4 or a ring of diamonds. We claim that G can be built from a 2-edge-connected cubic multigraph as in (iii). Suppose that G is a counter example with the minimum number of vertices.

If G has no diamonds, then every vertex of G is in exactly one triangle and therefore $V(G)$ can be partitioned into disjoint triangles. By contracting each triangle, we obtain a 2-edge-connected cubic multigraph H .

So G must have a string of diamonds. Let D be the set of vertices in the string of diamonds. Since G is cubic, G has two vertices not in D , say u and v , adjacent to D . If $u = v$, then because the degree of u is 3, u must have another incident edge e but e will be a cutedge of G . Thus $u \neq v$.

If u and v are adjacent in G , then u and v must have a common neighbor x , because otherwise G will have an induced subgraph isomorphic to $K_{1,3}$. However one of the edges incident with x will be a cutedge of G , a contradiction.

Thus u and v are nonadjacent in G . Let $G' = (G \setminus D) + uv$, that is obtained from G by deleting D and adding an edge uv . Then G' has no parallel edges or loops and moreover G' is 2-edge-connected claw-free cubic. Since G has a vertex not in a diamond, so does G' and therefore G' can be built from a 2-edge-connected cubic multigraph H by replacing some edges with strings of diamonds and replacing each vertex of H with a triangle. Since D is chosen maximally, u and v are not in diamonds and therefore H has the edge uv . So we can obtain G from H by doing all replacements to obtain G' and then replacing the edge uv with a string of diamonds. This completes the proof. \square

We remark that Proposition 1 can be seen as a corollary of the structure theorem of quasi-line graphs by Chudnovsky and Seymour [1]. A graph is a *quasi-line* graph if the neighborhood of each vertex is expressible as the union of two cliques. It is obvious that every claw-free cubic graph is a quasi-line graph. Chudnovsky and Seymour [1]

proved that every connected quasi-line graph is either a fuzzy circular interval graph or a composition of fuzzy linear interval strips. For 2-edge-connected claw-free cubic graphs, a fuzzy circular interval graph corresponds to a ring of diamonds and a composition of fuzzy linear interval strips corresponds to the construction (iii) of Proposition 1.

3 Main theorem

Theorem 2. *Every claw-free cubic n -vertex graph with no cutedge has more than $2^{n/12}$ perfect matchings.*

Proof. Let G be a claw-free cubic n -vertex graph with no cutedge. We may assume that G is connected. If G is isomorphic to K_4 , then the claim is clearly true. If G is a ring of diamonds, then G has $2^{n/4} + 1$ perfect matchings. Thus we may assume that G is obtained from a 2-edge-connected cubic multigraph H by replacing some edges of H with strings of diamonds and replacing each vertex of H with a triangle.

Let $k = |V(H)|$. In other words, $3k$ is the number of vertices not in a diamond of G . Equivalently, $V(G)$ can be partitioned into $(n - 3k)/4$ diamonds and k triangles each of which has exactly three distinct neighbors outside of the triangle.

Suppose that $k \geq n/6$. Since H has $3k/2$ edges, the cycle space of H has dimension $3k/2 - k + 1 = k/2 + 1$ and therefore $|\mathcal{C}(H)| = 2^{k/2+1}$. To obtain a 2-factor from $C \in \mathcal{C}(H)$, we transform C into a member $C' \in \mathcal{C}(G)$ so that it meets all 3 vertices of G corresponding to v for each vertex v of H incident with C as well as it meets all the vertices in each diamond that corresponds to an edge in C . Then for each vertex w of G unused yet in C' , we add a cycle of length 3 or 4 depending on whether the vertex is in a diamond; see Figure 2. Then this is a 2-factor of G because it meets every vertex of G . Since the complement of the edge-set of a 2-factor is a perfect matching, we conclude that G has at least $2^{k/2+1} \geq 2^{n/12+1}$ perfect matchings.

Now let us assume that $k < n/6$. We know that G has $(n - 3k)/4$ diamonds. The *length* of an edge e of H is the number of diamonds in the string of diamonds replaced with e . (If the edge e is not replaced with a string of diamonds, then the length of e is 0.)

Edmonds' characterization of the perfect matching polytope [4] implies that there exist a positive integer t depending on H and a list of $3t$ perfect matchings M_1, M_2, \dots, M_{3t} in H such that every edge of H is in exactly t of the perfect matchings. (In other words, H is fractionally 3-edge-colorable.) By taking complements, we have a list of $3t$ 2-factors of H such that each edge of H is in exactly $2t$ of the 2-factors in the list. Since G has $(n - 3k)/4$ diamonds, the sum of the length of all edges of H is $(n - 3k)/4$. Therefore there exists a 2-factor C of H whose length is at least $\frac{n-3k}{4} \frac{2}{3} = (n - 3k)/6$.

We claim that G has at least $2^{(n-3k)/6}$ 2-factors corresponding to C . For each diamond in the string replacing an edge e of C , there are two ways to route cycles of C through the diamond, see Figure 2. Since C passes through at least $(n - 3k)/6$ diamonds, G has at least $2^{(n-3k)/6}$ 2-factors. Since $k < n/6$, G has more than $2^{n/12}$ 2-factors. Thus G has more than $2^{n/12}$ perfect matchings. \square




Figure 2: Transforming a member of $\mathcal{C}(H)$ into a 2-factor of G (Solid edges represent edges in a member of $\mathcal{C}(H)$ or a 2-factor of G .)

We remark that every 3-edge-connected claw-free cubic n -vertex graph G has exactly $2^{n/6+1}$ perfect matchings, unless G is isomorphic to K_4 . That is because G has no diamonds and so, from the idea of the above proof, there is a one-to-one correspondence between the set of all 2-factors of G and the cycle space of a multigraph H obtained by contracting each triangle of G .

Note added in proof

A proof of the Lovász-Plummer conjecture has recently been submitted to the arXiv [5].

References

- [1] M. Chudnovsky and P. Seymour. The structure of claw-free graphs. In *Surveys in combinatorics 2005*, London Math. Soc. Lecture Note Ser. 327, pages 153–171. Cambridge Univ. Press, Cambridge, 2005.
- [2] M. Chudnovsky and P. Seymour. Perfect matchings in planar cubic graphs. Submitted, 2008.
- [3] R. Diestel. *Graph theory*, Graduate Texts in Mathematics 173. Springer-Verlag, Berlin, third edition, 2005.
- [4] J. Edmonds. Maximum matching and a polyhedron with 0, 1-vertices. *J. Res. Nat. Bur. Standards Sect. B*, 69B:125–130, 1965.
- [5] L. Esperet, F. Kardoš, A. King, D. Král' and S. Norine, Exponentially many perfect matchings in cubic graphs, arXiv:1012.2878v1, <http://arxiv.org/abs/1012.2878>.
- [6] L. Esperet, F. Kardoš, and D. Král'. Cubic bridgeless graphs have more than a linear number of perfect matchings. Accepted to Eurocomb'09, 2009.
- [7] M. Las Vergnas. A note on matchings in graphs. *Cahiers Centre Études Recherche Opér.*, 17(2-3-4):257–260, 1975. Colloque sur la Théorie des Graphes (Paris, 1974).
- [8] L. Lovász and M. D. Plummer. *Matching theory*, North-Holland Mathematics Studies 121. North-Holland, Amsterdam, 1986. Annals of Discrete Mathematics, 29.
- [9] E. M. Palmer, R. C. Read, and R. W. Robinson. Counting claw-free cubic graphs. *SIAM J. Discrete Math.*, 16(1):65–73, 2002.

- [10] J. Petersen. Die Theorie der regulären graphs. *Acta Math.*, 15(1):193–220, 1891.
- [11] D. P. Sumner. Graphs with 1-factors. *Proc. Amer. Math. Soc.*, 42:8–12, 1974.
- [12] M. Voorhoeve. A lower bound for the permanents of certain $(0, 1)$ -matrices. *Nederl. Akad. Wetensch. Indag. Math.*, 41(1):83–86, 1979.